

Subpart B—Specific New Animal Drugs for Use in Animal Feeds**§ 558.55 Amprolium.**

(a) *Approvals.* Type A medicated articles: 25 percent to No. 016592 in § 510.600(c) of this chapter for use as in paragraph (d) of this section.

(b) *Special considerations.* Do not use in Type B or Type C medicated feeds containing bentonite.

(c) *Related tolerances.* See § 556.50 of this chapter.

(d) *Conditions of use—(1) Cattle.* It is used as follows:

Amprolium in grams per ton	Indications for use	Limitations	Sponsor
(i) 113.5 to 11, 350; to provide 5 milligrams per kilogram of body weight per day.	Calves: As an aid in the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>E. zuernii</i> .	Top-dress on or mix in the daily ration. Feed for 21 days when experience indicates that coccidiosis is likely to be a hazard, as the sole source of amprolium. Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.	016592
(ii) 113.5 to 11, 350; to provide 10 milligrams per kilogram of body weight per day.	Calves: As an aid in the treatment of coccidiosis caused by <i>Eimeria bovis</i> and <i>E. zuernii</i> .	Top-dress on or mix in the daily ration. Feed for 5 days as the sole source of amprolium. Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.	016592

(2) *Chickens.* It is used as follows:

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 36.3 to 113.5	Replacement chickens: For development of active immunity to coccidiosis.	Feed continuously until onset of production as follows:	016592

Growing conditions	Up to 5 weeks of age	From 5 to 8 weeks of age	Over 8 weeks of age
	Amprolium in grams per ton	Amprolium in grams per ton	Amprolium in grams per ton
Severe exposure to coccidiosis	113.5 (0.0125%)	72.6–113.5 (0.008%–0.0125%)	36.3–113.5 (0.004%–0.0125%)
Moderate exposure to coccidiosis	72.6–113.5 (0.008%–0.0125%)	54.5–113.5 (0.006%–0.0125%)	36.3–113.5 (0.004%–0.0125%)
Slight exposure to coccidiosis	36.3–113.5 (0.004%–0.0125%)	36.3–113.5 (0.004%–0.0125%)	36.3–113.5 (0.004%–0.0125%)

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(ii) 36.3 to 113.5	Bacitracin methylenedisalicylate 4 to 50.	Replacement chickens: For development of active immunity to coccidiosis; and for increased rate of weight gain and improved feed efficiency.	Feed according to subtable in item (i). Bacitracin methylenedisalicylate as provided by No. 054771 in § 510.600(c) of this chapter.	054771
(iii) 72.6 to 113.5	Broiler chickens: For prevention of coccidiosis caused by <i>Eimeria tenella</i> only.	Feed continuously as the sole ration; as sole source of amprolium.	016592
(iv) 72.6 to 113.5	Bambermycins 1 to 2.	Broiler chickens: For prevention of coccidiosis caused by <i>Eimeria tenella</i> only; and for increased rate of weight gain and improved feed efficiency.	Feed continuously as the sole ration; as sole source of amprolium. Bambermycins as provided by No. 016592 in § 510.600(c) of this chapter.	016592
(v) 113.5	1. Laying chickens: For prevention of coccidiosis.	Feed continuously as the sole ration; as the sole source of amprolium.	016592

§ 558.58

21 CFR Ch. I (4–1–16 Edition)

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(vi) 113.5 to 227	2. Laying chickens: For treatment of coccidiosis in moderate outbreaks. 1. Replacement chickens: For prevention of coccidiosis where immunity to coccidiosis is not desired.	Feed for 2 weeks. Feed continuously from day-old until onset of production; as the sole source of amprolium.	016592
(vii) 113.5 to 227	Bambermycins 1 to 2.	2. Broiler chickens: For prevention of coccidiosis where immunity to coccidiosis is not desired. Broiler chickens: For prevention of coccidiosis where immunity to coccidiosis is not desired; and for increased rate of weight gain and improved feed efficiency.	Feed continuously as the sole ration; as sole source of amprolium. Feed continuously as the sole ration; as sole source of amprolium. Bambermycins as provided by No. 016592 in § 510.600(c) of this chapter.	016592
(viii) 227	Laying chickens: For treatment of coccidiosis in severe outbreaks..	Feed for 2 weeks	016592

(3) *Turkeys*. It is used as follows:

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 113.5	Bambermycins 1 to 4.	Growing turkeys: For prevention of coccidiosis; and for increased rate of weight gain and improved feed efficiency.	Feed continuously as the sole source of amprolium; bambermycins as provided by No. 016592 in § 510.600(c) of this chapter.	016592
(ii) 113.5 to 227	Turkeys: For prevention of coccidiosis.	Feed continuously as the sole ration; as sole source of amprolium.	016592

(4) *Pheasants*. It is used as follows:

Amprolium in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 159	Growing pheasants: For the prevention of coccidiosis caused by <i>Eimeria colchici</i> , <i>E. duodenalis</i> , and <i>E. phasiani</i> .	Feed continuously as sole ration. Use as sole source of amprolium.	016592
(ii) [Reserved]				

[41 FR 10985, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.55, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and at www.fdsys.gov.

§ 558.58 Amprolium and ethopabate.

(a) *Specifications*. Type A medicated articles containing:

(1) 25 percent amprolium and 8 percent ethopabate or 5 percent amprolium and 1.6 percent ethopabate;

(2) 25 percent amprolium and 0.8 percent ethopabate or 5 percent amprolium and 0.16 percent ethopabate.

(b) *Approvals*. See No. 016592 in § 510.600(c) of this chapter.

(c) *Special considerations*. Do not use in Type B or Type C medicated feeds containing bentonite.

(d) *Related tolerances*. See §§ 556.50 and 556.260 of this chapter.

(e) *Conditions of use*. It is used in chicken feed as follows: